

Kinematic Evidence of Minor Mergers in Normal Sa Galaxies: NGC 3626, NGC 3900, NGC 4772 and NGC 5854

Martha P. Haynes^{1,2}

Center for Radiophysics and Space Research and National Astronomy and Ionosphere Center³, Cornell University, Space Sciences Building, Ithaca NY 14853
haynes@astrosun.tn.cornell.edu

Katherine P. Jore^{1,2}

Department of Physics and Astronomy, University of Wisconsin at Stevens Point, Stevens Point, WI 54481
kjore@uwsp.edu

Elizabeth A. Barrett

Department of Astronomy and Center for Radiophysics and Space Research, Cornell University, Space Sciences Building, Ithaca NY 14853
eab15@cornell.edu

Adrick H. Broeils^{1,4}

Center for Radiophysics and Space Research, Cornell University, Space Sciences Building, Ithaca NY 14853
adrick.broeils@nl.origin-it.com

Brian M. Murray

Linfield College, 900 SE Baker St., McMinnville, OR 97128
bmurray@linfield.edu

ABSTRACT

BVRI and H α imaging and long-slit optical spectroscopic data are presented for four morphologically normal and relatively isolated Sa galaxies, NGC 3626, NGC 3900, NGC 4772 and NGC 5854. VLA H I synthesis imaging is presented for the first three objects. In all four galaxies, evidence of kinematic decoupling of ionized gas components is found in the long-slit spectroscopic data; the degree and circumstances of the

¹Visiting Astronomer, National Radio Astronomy Observatory which is a facility of the National Science Foundation operated under a cooperative agreement by Associated Universities, Inc.

²Visiting Astronomer, Kitt Peak National Observatory which is part of the National Optical Astronomy Observatories, operated by the Association of Universities for Research in Astronomy, Inc. (AURA) under a cooperative agreement with the National Science Foundation.

³The National Astronomy and Ionosphere Center is operated by Cornell University under a cooperative agreement with the National Science Foundation.

⁴Now at Origin Nederland B.V., Nieuwegein, The Netherlands.

distinct kinematics vary from complete counterrotation of all of the gas from all of the stars (NGC 3626) to nuclear gas disks decoupled from the stars (NGC 5854) to anomalous velocity central gas components (NGC 3900 and NGC 4772). In the three objects mapped in H I, the neutral gas extends far beyond the optical radius, $R_{HI}/R_{25} \geq 2$. In general, the H I surface density is very low and the outer H I is patchy and asymmetric (NGC 3900) or found in a distinct ring, exterior to the optical edge (NGC 3626 and NGC 4772). While the overall H I velocity fields are dominated by circular motions, strong warps are suggested in the outer regions by bending of the minor axis isovelocity contours (NGC 3900) and/or systematic shifts in position angle between inner and outer rings (NGC 3626 and NGC 4772). In the interior, coincidence is found between the H α and H I in rings, sometimes partial and crisscrossed by dustlanes. Optical imaging is also presented for NGC 4138 previously reported by Jore *et al.* (1996) to show counterrotating stellar components. The multiwavelength evidence is interpreted in terms of the kinematic “memory” of past minor mergers in objects that otherwise exhibit no morphological signs of interaction.

Key words: – galaxies: evolution – galaxies: individual (NGC 3626, NGC 3900, NGC 4138, NGC 4772, NGC 5854) – galaxies: interactions – galaxies: kinematics and dynamics – galaxies: spiral

1. Introduction

While mergers between comparable mass galaxies may be responsible for some of the most dramatic extragalactic events, minor mergers with small mass satellites may play a subtle but nonetheless critical role in the evolution of garden-variety spirals, including perhaps the Milky Way. It is easy to recognize the fireworks of the former by the tides, starbursts and nuclear activity they trigger. The latter however may lurk hidden to morphological inspection. Minor mergers are suggested to produce the X-shaped structures seen in some peculiar S0's (Mihos *et al.* 1995), the departures from axisymmetry seen in a significant fraction of spiral disks (Zaritsky & Rix 1997), the driving mechanism for strong stellar bars (Laine & Heller 1999) and the peculiar extended counter-rotating disks seen in some early-type spirals, e.g. NGC 7217 (Merrifield & Kuijken 1994; Buta *et al.* 1995), NGC 3593 (Bertola *et al.* 1996; Corsini *et al.* 1998), NGC 3626 (Ciri *et al.* 1995) and NGC 4138 (Jore *et al.* 1996).

Satellites are commonly found in the vicinity of normal galaxies (Zaritsky *et al.* 1997). If the Local Group is at all typical of spiral extragalactic environments, then the characteristics of such satellites are familiar. Interaction of the Milky Way with its low mass satellites is hypothesized to explain a variety of local kinematic peculiarities: the thick disk, the outer flare and warp, the Magellanic Stream and the high velocity H I clouds.

Several authors have begun to address numerically the complex phenomenon of minor mergers of gas-rich satellites with more massive spiral companions. An immediate worry is how to avoid over-heating the inner disk of the primary or disrupting it entirely. Minor mergers of low-density, small mass, gas-rich companions may avoid vertical heating, thereby preserving the victim disk. Predictions of the observable consequences of the accretion of a small mass companion by a spiral include the warping or actual spreading out of the disk (Quinn *et al.* 1993). As the satellite material sinks towards the center of the large galaxy, the gas, unlike the stars, may lose angular momentum because of dissipation (Hernquist & Mihos 1995). The resultant inflow of gas towards the center may trigger a nuclear starburst (Mihos & Hernquist 1994) or fuel nuclear activity (Taniguchi 1999). Because of the predicted heating, even a satellite with one tenth the mass will thicken the disk and possibly build-up the

bulge, driving the morphology of the post merger primary towards an earlier spiral type. The discovery of kinematically decoupled disks in early spirals may thus prove the efficacy of minor merger models.

Intrinsic galaxy properties arise from two principal components: absolute “scale” (size, luminosity, mass) and “form” (morphology, bulge-to-disk ratio, color, gas content, star formation rate). It is well-known (e.g. Roberts & Haynes 1994) that the early spiral types show a much greater spread in their characteristic properties related to “form” than do their later-type counterparts. As a part of a study to investigate the heterogeneity and dark matter content of the Sa galaxy class, we have investigated the kinematics of gas and stars in a sample of morphologically normal, isolated galaxies classified as Sa in the *Revised Shapley Ames Catalog* (Sandage & Tammann 1987; RSA). In the course of that study, we have obtained major and minor axis long slit spectra for both stars and gas for a sample of 20 objects; for nine of them, we have also obtained H I synthesis maps yielding also two-dimensional H I velocity fields. A summary of results focussing on the data presentation and mass modelling was presented in Jore (1997). The analysis of the mass modelling and stellar velocities will be presented elsewhere (Jore *et al.*, in preparation).

During the course of that work, a number of unusual cases of kinematic decoupling were discovered. In Jore *et al.* (1996), we discussed the distinctive case of NGC 4138, an isolated Sa with two extended coplanar counter-rotating stellar disks embedded in a huge H I disk that co-rotates with the secondary stellar component. In a separate paper (Jore *et al.* 2000), we discuss five Sa galaxies, NGC 1169, 3623, 4866, 5377 and 5448, which represent a range of environments and morphologies within the Sa class. Of those objects, NGC 1169 and NGC 5448, exhibit no peculiar optical kinematics but their outer H I disks are strongly warped and kinematically skewed. NGC 3623, whose companions in the Leo Triplet are known to be interacting, exhibits a truncated H I disk and central gas decoupling. NGC 4866, viewed nearly edge-on, and NGC 5377 show doubling of their central gas components, as well as outer H I warps. Here, we present four additional examples: NGC 3626, NGC 3900, NGC 4772 and NGC 5854. Along with NGC 4138, each of these morphologically normal Sa galaxies exhibits peculiar kinematics that betray a disturbed past. We propose that each represents a different stage and circumstance of involve-

ment in a relatively minor merger event with a gas-rich companion that is no longer recognizable.

This paper presents evidence of this past interaction through a combination of optical imaging and spectroscopy and H I synthesis mapping. In Section 2, we discuss the acquisition and reduction of the body of data: broad- and narrow-band optical images, optical long-slit spectra, and H I synthesis maps. The kinematic evidence of past interaction is presented for each galaxy in Section 3. Section 4 gives a discussion of the observational results in the context of the minor merger scenario.

2. Observations

The original observing program was designed to provide the details of the galaxy dynamics for the purpose of modelling the dark matter distribution in Sa galaxies and, as such, consisted of long-slit major and minor axis spectra of both stars and ionized gas. For a subset, principally gas-rich objects showing evidence in the optical spectra of kinematic peculiarities, H I synthesis mapping and photometric broad- and narrow-band imaging was also undertaken.

The basic properties of these four galaxies plus NGC 4138 discussed by Jore *et al.* (1996) are summarized in Table 1. The first entries include the (B1950) R.A. and Dec. and the morphological classifications as given in both the RSA (Sandage & Tammann 1987) and the *Third Reference Catalog of Bright Galaxies* (de Vaucouleurs *et al.* 1991; RC3). A morphological description of the H α emission is then given, followed by a set of radial measures: $D_{25} \times d_{25}$, in arcminutes, from the RC3 and the corresponding radius R_{25} , in arcseconds; R_{B25} , in arcseconds, derived from the B image; R_{I83L} , in arcseconds, the radius encompassing 83% of the total I band light, following Haynes *et al.* (1999a); and R_{orc} , in arcseconds, the maximum radius at which the optical rotation curve is derived, for any species, stars or gas. The corrected integrated blue magnitude, B_T^0 , and the color indices, $(U-B)$ and $(B-V)$, all from the RC3, are then listed, along with the optical heliocentric velocity, $V_\odot$, in km s^{-1} , and the adopted distance, in Mpc. Since all of these galaxies are nearby, distances are adopted from available estimates accounting for group membership and local peculiar motions and/or supernova distance moduli. The total B luminosity, L_B , is derived from B_T^0 and the distance. The total apparent magnitudes m_B , m_V , m_R , and m_I , and their estimated errors,

have been derived from the images following Haynes *et al.* (1999a). Likewise, the ellipticity $\epsilon = (1 - \frac{b}{a})$ and position angle are the mean disk properties extracted from the I band isophotal fits using the “marking the disk” method of Haynes *et al.* (1999a). Finally, the adopted R band bulge-to-disk ratio B/D_R is given.

The acquisition and reduction of the multiwavelength data sets are discussed here separately.

2.1. Optical Images

BVR and H α images were obtained with the KPNO 0.9m telescope in its f/7.5 configuration over the nights of April 19–23, 1996. Additionally, I band images, some taken under non-photometric conditions, were obtained with the same instrument by R. Giovanelli and MPH as part of another project. For all images, the 2048×2048 pixel T2KA CCD was used. Its read noise is 4 e^- and the gain was set to $5.4 \text{ e}^- \text{ ADU}^{-1}$, except for the I band images, when the gain was set to $3.6 \text{ e}^- \text{ ADU}^{-1}$. With the field corrector in, the images had a pixel scale of $0.68''$. In this section, we describe the image acquisition, processing and analysis techniques.

2.1.1. Broadband BVRI Images

The broadband images were exposed over the whole chip, giving a total field of view of $23'$ on a side. The images were trimmed, bias subtracted, and flat fielded using CCDPROC in IRAF⁵. Flat fields for B , V and R were obtained from twilight sky observations; for the I band, a flat field was constructed from the median of 30–50 object frames taken throughout the night. Further image reductions made use of the GALPHOT⁶ package of IRAF/STSDAS scripts for galaxy surface photometry (see Haynes *et al.* 1999a). Regions of the image free of galaxy emission and bright stars were interactively marked. An average value of the sky contribution in these boxes after masking of the stars within them was subtracted as the sky contribution. Images were also edited interactively to remove stars and cosmic rays and convolved to guarantee the same resolution before combining. Images with three or more observations on

⁵IRAF is distributed by the National Optical Astronomy Observatories.

⁶The GALPHOT surface photometry package is a collection of IRAF/STSDAS scripts originally developed by Wolfram Freudling and John Salzer; the Cornell version has been further modified and is maintained by MPH.

TABLE 1
GALAXY PROPERTIES

	N 3626 U 6343	N 3900 U 6786	N 4138 U 7139	N 4772 U 8021	N 5854 U 9726
R.A.(B1950)	111725.9	114633.3	120659.3	125055.9	150516.2
Dec.(B1950)	+183756	+271806	+435757	+022627	+024537
RSA Type ^a	Sa	S(r)a	S(r)a pec	Sa:	Sa
RC3 Type ^b	RLAT+..	.LAR+..	.LAR+..	.SAS1..	LBS+./
H α morph ^c	n+cr	(trace)	n+cb+or	n+cb+or	...
$D_{25} \times d_{25}$ ^b	2.69x 1.95	3.16 x 1.70	2.57 x 1.70	3.39 x 1.70	2.75 x 0.79
R_{25} (") ^b	81	96	78	102	84
R_{B25} (")	76	97	85	132	78
R_{I83L} (")	56	75	54	120	48
R_{orc} (") ^d	45	53	50	58	51
B_T^0 ^b	11.75	12.24	12.14	11.90	12.68
$(U - B)$ ^b	0.30	...	0.31	0.26	0.29
$(B - V)$ ^b	0.81	0.82	0.83	0.83	0.81
$V_{\odot}$ (km s ⁻¹) ^b	1493	1798	888	1040	1663
Distance (Mpc)	24	27	16	16	24
L_B ($10^{10} L_{\odot}$)	1.8	1.4	0.6	0.7	0.8
m_B	11.91 $\pm$.04	12.40 $\pm$.02	12.12 $\pm$.04	11.97 $\pm$.04	...
m_V	11.11 $\pm$.02	11.65 $\pm$.02	11.31 $\pm$.02	11.15 $\pm$.03	11.20 $\pm$.10
m_R	10.59 $\pm$.02	11.08 $\pm$.02	10.73 $\pm$.03	10.58 $\pm$.02	10.64 $\pm$.02
m_I	...	10.42 $\pm$.02	10.03 $\pm$.02	...	...
ϵ	0.32 $\pm$ 0.02	0.48 $\pm$ 0.02	0.39 $\pm$ 0.02	0.48 $\pm$ 0.03	0.67 $\pm$ 0.02
P.A. ($^{\circ}$)	164 $\pm$ 3	4 $\pm$ 1	150 $\pm$ 1	148 $\pm$ 2	55 $\pm$ 1
B/ D_R	0.37	0.27	0.19	0.44	0.82

^aFrom RSA.

^bFrom RC3.

^cCode for morphology: n=nuclear; cr=core ring; cb=elongated core; or=outer ring.

^dMaximum extent of optical rotation curve traced by any species.

the same filter were scaled and median combined to remove cosmic rays. For images with fewer exposures cosmic rays were removed before combining using the task COSMIC in IRAF and interactively editing all remaining cosmic rays by hand. The filters, total exposure times, and the seeing values for the combined images are presented in Table 2 for each galaxy.

Elliptical isophotal fitting routines were run on the star edited *BVRI* images of each galaxy. These routines require the user to mark an initial major and minor axis. The program then fits isophotal ellipses moving outwards and inwards from the marked initial location and integrates the flux within the ellipses. Using these results, disk parameters such as position angle, ellipticity, and slope are fit and extrapolated magnitudes are found. Selected values of these parameters are included in Table 1.

Conversion from instrumental to apparent magnitudes was accomplished using frequent observations of standard stars (Landolt 1992) observed throughout each night. The photometric coefficients for two photometric nights during which the *BVR* images were obtained (nights 1 and 3) are listed in Table 3. Photometric coefficients were also determined for night 2; however, they fail to reproduce Landolt standard magnitudes accurately and were therefore discarded as non-photometric. All images from night 2 were subsequently scaled to ones obtained on nights 1 or 3 in order to insure photometric accuracy.

Color plots of each galaxy in $B - V$, $B - R$ and $V - R$ were produced by first aligning all the bandpasses and convolving them to the same resolution (the poorest of the set, as given in Table 2). The images were then converted into instrumental magnitudes and rescaled to apparent magnitudes using the appropriate photometric coefficients. Final color images were made by subtracting the resultant images and correcting for galactic extinction. Values for galactic extinction were calculated using $A_g(B)$ values from the RC3 and the conversions to other bands given in Rieke & Lebofsky (1985). Additionally, model color images were also created from the broadband isophotal ellipse fitting routines mentioned above. Using the results of the ellipse fitting, model broadband images were created, converted into apparent magnitudes as above, and subtracted to form noiseless model $B - V$, $B - R$ and $V - R$ images.

The separation of bulge and disk components was accomplished using FITMAP, a two-dimensional decomposition routine written and kindly provided by

G. Moriondo (Moriondo *et al.* 1998). FITMAP fits a model with a specified bulge and exponential disk to the data, adjusting the allowed parameters until the optimum model is found as judged by the reduced χ^2 value. The bulge shape index n for the generalized exponential law $I_b(R) = I_e \exp[-(R/R_e)^{1/n}]$ can be specified as an integer value, with $n = 1$, an exponential bulge and $n = 4$, a de Vaucouleurs bulge. In order to speed processing time, the fit was performed only on an averaged quadrant of the galaxy. In this data set, the average was typically a combination of all four quadrants, achieved by folding the image along both major and minor axes, except where bright stars were found close to the galaxy center. All filters were fitted separately to allow comparison of the results obtained with different bandpasses.

Recent literature (e.g. de Jong 1996, Moriondo *et al.* 1998) shows that for earlier type spirals, such as those used in this sample, an $n = 2$ bulge is the best fit. That finding is supported likewise here. Furthermore, the results indicate an increase in the B/D parameter with redder wavelengths, compatible with the bulge having an overall redder color than the disk and there being less obscuration from dust at longer wavelengths. A summary of the B/D parameters obtained for each filter and adopting the $n = 2$ bulge is given in Table 4.

2.1.2. Narrowband $H\alpha$ Images

$H\alpha$ images were produced by observing the same region in two narrow (80Å) bandpasses, one centered on the appropriately redshifted $H\alpha$ emission line (the ON image) and a second just off the emission line (the OFF image). Both OFF and ON images were obtained over the nights of April 22–23, 1996 with the KPNO 0.9m telescope. The same T2KA chip, as described above, was used; however, only the central 1024x1024 pixel region was exposed. Usually, two ON exposures of 30 minutes each and a single OFF one of the same duration were obtained; Table 2 gives the details of the $H\alpha$ observations for each galaxy. Images were trimmed, bias subtracted, flat fielded, and cosmic ray edited in the same manner as the *BVR* images described above. The ON and OFF images were then convolved to the same resolution, and the two ON images were combined. The OFF image was scaled to the summed ON image and subtracted to produce an $H\alpha$ line image. The nights were both non-photometric and no calibration is available.

TABLE 2
DETAILS OF IMAGING OBSERVATIONS

Object	Filter	Night	T_{exp} (sec)	Seeing "	Note
N3626	B	1	3600	2.69	a
	V	1	1350	2.40	a
	R	1	1080	1.97	
	I		600	2.99	b
	ON	4	3600	1.98	b
	OFF	4	1800	1.98	b
N3900	B	3	3600	2.03	
	V	3	1350	2.02	
	R	3	900	2.16	
	I		600	1.62	
	ON	5	3600	1.81	b
	OFF	4	1800	1.77	b
N4138	B	1	3600	2.24	
	V	1	1350	1.86	
	R	1	900	1.81	
	I		600	2.12	
	ON	5	3600	2.00	b
	OFF	4	1800	2.09	b
N4772	B	1	3600	1.57	
	V	3	1350	1.65	
	R	3	900	1.65	
	I		600	1.92	b
	ON	5	3600	1.56	b
	OFF	5	1800	1.64	b
N5854	B	1	3600	2.50	b
	V	1	1350	2.19	
	R	1	900	1.99	
	I		600	2.58	b

^aTracking problems; image quality poor

^bNonphotometric

TABLE 3
PHOTOMETRIC SOLUTIONS FOR BVR IMAGING

Night	Filter	Zeropoint (mag)	Color Coef.	Extinc. Coef.
1	B	21.005 (0.039)	0.099 (0.019)	0.237 (0.004)
	V	21.311 (0.018)	0.000 (0.009)	0.157 (0.004)
	R	21.411 (0.013)	0.009 (0.014)	0.104 (0.003)
3	B	21.062 (0.007)	0.075 (0.012)	0.233 (0.004)
	V	21.354 (0.013)	-0.017 (0.010)	0.155 (0.003)
	R	21.424 (0.010)	0.019 (0.016)	0.102 (0.003)

TABLE 4
RESULTS OF 2D BULGE–DISK DECOMPOSITION.

Object	Filter	Exponential Disk		$n = 2$ Bulge		χ^2	B/D	
		μ_o (mag/arcsec ²)	h (arcsec)	μ_e (mag/arcsec ²)	R_e (arcsec)			
N3626	B	20.94(0.01)	21.3(0.16)	18.94(0.05)	3.2(0.08)	2.73	0.37	
	V	20.10(0.01)	21.1(0.12)	17.99(0.05)	3.0(0.06)	3.75	0.37	
	R	19.55(0.01)	20.7(0.13)	17.50(0.04)	3.0(0.06)	3.20	0.37	
	I	...	21.3(0.34)	...	3.4(0.05)	1.16	0.30	a
N3900	B	21.68(0.02)	24.4(0.21)	20.69(0.05)	4.8(0.18)	1.55	0.25	
	V	20.78(0.01)	23.0(0.16)	19.77(0.04)	4.9(0.13)	1.61	0.30	
	R	20.15(0.01)	22.2(0.14)	19.16(0.04)	4.8(0.12)	1.84	0.27	
	I	19.27(0.01)	20.7(0.11)	18.25(0.02)	4.0(0.06)	0.79	0.25	
N4138	B	20.46(0.03)	16.6(0.14)	21.44(0.08)	7.0(0.48)	2.19	0.19	
	V	19.67(0.01)	16.9(0.08)	19.90(0.04)	5.2(0.16)	1.64	0.20	
	R	19.06(0.01)	16.7(0.06)	19.01(0.04)	4.4(0.11)	1.72	0.19	
	I	18.37(0.01)	17.0(0.05)	17.95(0.02)	3.7(0.06)	1.02	0.18	
N4772	B	22.81(0.01)	50.4(0.36)	21.22(0.01)	9.6(0.10)	0.44	0.41	
	V	21.96(0.01)	48.4(0.25)	20.27(0.01)	9.3(0.08)	0.65	0.45	
	R	21.35(0.01)	46.8(0.23)	19.64(0.01)	9.1(0.07)	0.72	0.47	
	I	...	42.3(0.40)	...	8.1(0.07)	0.23	0.44	a
N5854	B	...	20.0(0.38)	...	6.0(0.14)	0.66	0.63	a
	V	20.15(0.22)	19.5(0.35)	19.07(0.23)	5.7(0.12)	0.55	0.61	a
	R	19.59(0.02)	19.5(0.15)	18.54(0.03)	5.7(0.12)	0.73	0.59	
	I	...	19.6(0.17)	...	5.6(0.11)	1.05	0.63	a

^aNonphotometric

2.2. Optical Long Slit Spectra

These data are described in more detail in Jore (1997). Optical long slit spectra were taken at the Palomar 5m Hale Telescope⁷ using the red and blue cameras of the Double Spectrograph. The slit dimensions were 120'' by 2''. A dichroic at 5500 Å and gratings in both cameras with dispersions of 1200 lines mm⁻¹ were used. The red camera was centered on the emission lines of H α and [N II] λ 6548 Å and λ 6584 Å, with the [S II] λ 6717 Å and λ 6731 Å lines also present. The blue camera was centered on the Mg I b λ 5183.6, 5172.7 and 5167.3 Å stellar absorption lines with the [O III] λ 5007 Å emission line also visible. Table 5 summarizes the Double Spectrograph setup, while Table 6 presents the details of all spectral observations used to extract rotation curves. All exposures over 40 minutes composed of co-added shorter exposures. It should be noted that some spectra were taken under conditions of poor transparency so that the effective integration is significantly less than the actual exposure time. Arc lamps and radial velocity standard stars were observed for wavelength and velocity calibration. In general, arc lamps were used for wavelength calibration on the blue side, while night sky lines provided the necessary scaling anchor on the red side.

Rotation curves were extracted for each of the ionized gas components using a Gaussian fitting routine. The stellar rotation curves were extracted using several different methods employing both cross-correlation and direct-fitting techniques. Further discussion of the reduction procedure is presented in Jore *et al.* (1996) and Jore (1997). The gas rotation curves displayed in Section 3 are derived from H α and/or [NII], depending on the relative strengths of the lines and the possible confusion with line absorption in the case of H α . The stellar rotation curves are discussed here; the stellar velocity dispersions will be discussed elsewhere (Jore *et al.*, in preparation).

2.3. HI Synthesis Imaging

The H I 21-cm line observations were conducted with the Very Large Array (VLA)⁸ in its C configuration. As discussed in Section 3.4, NGC 5854 is very

gas-poor and was not mapped. A summary of the observational setup for the VLA observations of the other three galaxies is given in Table 7. Each galaxy was allocated 8 hours of array time in its C configuration, with a nominal beam size of $\sim 12''$ and sensitivity to structures smaller than 7'. Within each 8 hour allocation, a flux calibrator was observed three times for ~ 10 minutes, and a phase calibrator was observed for 4 minutes between each 50 minute observation on the galaxy. All observations were obtained using the 4ABCD spectral line mode with two overlapping spectral bands of 32 channels per band, with an overlap of at least 10 channels so that the total number of channels was 54, each 20.6 km s⁻¹ wide. The total amount of on-source integration time and the total velocity coverage of the combined spectral bands are listed in Table 7.

The UV data were flux and phase calibrated using the AIPS software package after flagging bad visibilities. The bandpass for each antenna of the line data was calibrated for each IF separately. The two spectral bands were then combined using the task UVGLU. Dirty maps were made using MX and were inspected to determine the channels free of line emission. The continuum was subtracted in the UV plane using UVLIN by interpolating the continuum from the emission-line free channels to those with line emission. Cleaned map cubes were created from the UV data using a Gaussian beam derived from the FWHM of the dirty beam. For most of the galaxies, two maps with different resolutions were made; one (using natural weighting) with lower spatial resolution and higher signal-to-noise, and the other (using uniform weighting) with higher resolution but noisier. The beam size and position angle and RMS noise per channel for each map are listed in Table 7.

Once the map cubes were made, further analysis of the data was done using the GIPSY (van der Hulst *et al.* 1992) data reduction package. The data cubes were first smoothed to twice the original beam size and then clipped at the 1 σ level. The regions containing line emission in two or more consecutive channels were interactively extracted. The data in the original cube corresponding to the regions extracted in the smoothed cube were conditionally transferred. This final cube contains only the line emission from the sources. Zeroth and first order moment maps (integrated intensity and velocity field maps) were made using the GIPSY task MOMENTS.

Column density maps were obtained from the in-

⁷Observations at the Palomar Observatory were made as part of a continuing collaborative agreement between the California Institute of Technology and Cornell University.

⁸The Very Large Array is a facility of the National Radio Astronomy Observatory.

TABLE 5
DOUBLE SPECTROGRAPH OBSERVING SETUP

Dates	Camera	Chip	Spat. Res. ($'' \text{ pix}^{-1}$)	λ Range ($\text{\AA}$)	λ Res. ($\text{\AA} \text{ pix}^{-1}$)	Vel. Res. (km s^{-1})
Mar. 94 -	Red	TI 167	0.58	6190-6850	0.82	38
Apr. 95	Blue	TI 432	0.78	4950-5400	0.56	38
Mar. 96	Red	1396CR14-0	0.46	6150-6825	0.66	30
	Blue	TI 432	0.78	4950-5400	0.56	38

TABLE 6
OPTICAL LONG-SLIT SPECTRAL OBSERVATIONS

Object	Date	Total Int. (seconds)	PA ($^{\circ}$ E of N)	Axis
NGC 3626	April 1995	3600	157	major
		1800	67	minor
NGC 3900	March 1994	3600	2	major
		3600	92	minor
	April 1995	3600	2	major
		1800	92	minor
NGC 4772	April 1995	3600	147	major
		1800	57	minor
NGC 5854	March 1994	3600	55	major
		3000	145	minor
	April 1995	3600	55	major
		1800	145	minor
	March 1996	6419	55	major

TABLE 7
VLA-C HI SYNTHESIS OBSERVING PARAMETERS

	weighting scheme	N 3626 U 6343	N 3900 U 6786	N 4772 U 8021
Date Obs.		2/23/96	2/24/96	11/10/94
T _{int} (min)		373.5	371.5	403
Velocity Coverage (km s ⁻¹)		930 – 2012	1251 – 2336	485 – 1564
Beam Size (")	uniform	12.1 x 11.8	11.7 x 11.7	13.8 x 12.4
	natural	16.8 x 16.2	22.5 x 20.0	31.7 x 23.1
Beam PA (°)	uniform	113	165	143
	natural	101	79	137
RMS channel ⁻¹ (mJy beam ⁻¹)	uniform	0.28	0.36	0.44
	natural	0.23	0.31	0.33
minimum N _{HI} (10 ²⁰ atoms cm ⁻²)	natural	1.0	0.3	0.2

egrated intensity maps using the relation

$$N(H) = 1.10297 \times 10^{24} (1+v/c)^2 \frac{1}{ab} \int S(v) dv \text{ [atoms cm}^{-2}] \quad (1)$$

where $\int S(v) dv$ is the integrated flux (in Jy/beam) times the velocity width (in km s⁻¹/channel), a and b are the FWHM of the major and minor axes of the beam (in arcsec), and v is the systemic velocity of the source (in km s⁻¹). The minimum detectable column density quoted in Table 7 is the 2σ value in the natural weight column density map.

2.4. Continuum emission

Cleaned maps were made of the non-continuum subtracted maps. The continuum map for each source was determined by taking the mean of all line-free channels. The primary beam corrected flux density of nearby sources was calculated using the GIPSY routine FLUX. The continuum results for each galaxy are discussed separately; the derived continuum flux density is included in Table 8.

2.4.1. NGC 3626

For NGC 3626, the average of 17 line-free channels was calculated, yielding a continuum map. Continuum emission was detected at the center of NGC 3626 with a flux density $S_{1413} = 9.9$ mJy, slightly larger than the flux density $S_{1415} = 6$ mJy measured by van Driel *et al.* (1989). There is another nearby source located 2'9 west of the center (11^h17^m13^s.5,

+18°37'38" (1950)) with a flux density $S_{1413} = 4.1$ mJy. The companion galaxy, UGC 6341, has no continuum emission associated with it, at a 3σ point source sensitivity of 0.3 mJy.

2.4.2. NGC 3900

For NGC 3900, the continuum map was constructed similarly. Continuum emission was detected at the center of NGC 3900 with a flux density $S_{1412} = 1.2$ mJy in agreement with the upper limit of $S_{1415} < 6$ mJy given by van Driel *et al.* (1989). A nearby source is located 2'8 north of the center (11^h46^m35^s.3, +27°20'24" (1950)) with a flux density $S_{1412} = 1.4$ mJy which is coincident with a small background galaxy seen in the optical images.

2.4.3. NGC 4772

For NGC 4772, the average of 16 line-free channels was used to produce the continuum map. Continuum emission was detected at the center of NGC 4772 with a flux density $S_{1415} = 3.4$ mJy. Four other nearby sources were found. Their positions and flux densities are as follows: (a): 12^h50^m59^s.9, +02°24'43" (1950) with a flux density $S_{1415} = 5.0$ mJy; (b): 12^h51^m06^s.0, +02°26'39" (1950) with a flux density $S_{1415} = 4.0$ mJy; (c): 12^h51^m09^s.5, +02°28'15" (1950) with a flux density $S_{1415} = 2.3$ mJy; and (d): 12^h50^m58^s.3, +02°29'35" (1950) with a flux density $S_{1415} = 3.7$ mJy.

2.5. HI Channel Maps and Global Profiles

The channels maps extracted from the natural weight data cubes are shown in Figures 1 to 3 for the individual galaxies NGC 3626, NGC 3900 and NGC 4772 respectively. The global HI profiles extracted for the three target galaxies and for two others, UGC 6341 and NPM1G+18.0219, both found in the field of NGC 3626, are shown in Figure 4. Table 8 presents the properties of these five objects as derived from the VLA maps; NGC 4138 is also included for comparative purposes. Entries are: (1) the continuum flux density derived from the VLA map as discussed in Section 2.4, in mJy; (2) the H I deficiency, $< DEF >$, given by Magri (1994). $< DEF >$ is the logarithmic difference between the observed H I mass and that expected for an isolated galaxy of the same morphological class and linear *optical* diameter (Haynes & Giovanelli 1984); (3) the H I line flux $\int S_{HI} dV$ in Jy-km s⁻¹. For the VLA datasets, $\int S_{HI} dV$ is derived from the natural weight data cube, after application of a primary beam correction. (4) the H I mass, derived from $\int S_{HI} dV$, in M_⊙; (5) the H I mass to blue luminosity ratio, M_{HI}/L_B , in solar units; (6) the H I line heliocentric systemic velocity, V_{21} , in km s⁻¹; (7) the width of the H I line global profile at 20% of the peak intensity, ΔV_{21} , in km s⁻¹; (8) the ratio of the H I to optical radius, R_{HI}/R_{25} , where R_{HI} is measured at a constant surface density of 1 M_⊙ pc⁻²; (9) the position of the dynamical center; (10) the inclination derived from the tilted ring model, in degrees; (11) the maximum rotational velocity derived from the best-fit rotation curve, V_{max} , in km s⁻¹. Further discussion is deferred to the next section in which each galaxy is discussed individually.

3. Results for Individual Objects

3.1. NGC 3626

NGC 3626 is found in an extended cloud of galaxies related to the NGC 3607 group, LGG 237 (Garcia 1993); the central concentration around NGC 3607 is located about 50' (350 kpc at the adopted distance of 24 Mpc) away. The only object known to be associated with the group that lies within 30' of NGC 3626 is UGC 6341, a Magellanic spiral at similar redshift, located at a separation of 5'.9, 41 kpc. UGC 6341 was also detected in the VLA H I synthesis map and is discussed further below. A ten minute exposure with the Palomar 5m telescope and the Double Spectrograph of the faint galaxy located at R.A.(1950) =

11^h17^m30^s.1, Dec.(1950) = +18°34'31" failed to reveal any lines and its redshift is otherwise unknown.

Analyzing long-slit spectra of both the stars and ionized gas, Ciri *et al.* (1995; see also Jore 1997) found that all of the gas in this galaxy is counterrotating relative to the stars. Recent CO observations by García-Burillo *et al.* (1998) reveal the presence of a massive molecular nuclear disk associated with the ionized gas. The H I has been mapped previously by van Driel *et al.* (1989) with the Westerbork Synthesis Radio Telescope (WSRT).

As evident in Figure 5, NGC 3626 is a ringed Sa galaxy with smooth, faint spiral structure and a prominent dustlane to the west of the nucleus. The three left panels of Figure 5 show different representations of the *R* band image, displayed with different contrast levels and scaling to emphasize the low surface brightness outer disk (top), the ring (center) and the dustlane (bottom). Unfortunately, the telescope suffered poor tracking when the *B* and *V* images were obtained but, while the PSF is degraded, colors can still be derived. The *B* - *V* color image shows no large scale structure in the disk aside from the dustlane. The reddening seen in both *B* - *V* and *B* - *R* at +10" along the minor axis coincides with the dustlane. The bulge is small, dominating the inner *R* < 9", with a *B* - *V* color of ~ 0.7; two larger blue (*B* - *V* ~ 0.4) components are found about 6" to the NW and SE of the bulge. As shown in the top right panel of Figure 5, the *R* band surface brightness profile is well fit by an $n = 2$ bulge, yielding a modest $B/D_R \sim 0.37$. Similar values are obtained for the other bands as given in Table 4. Structure seen in the surface brightness profile coincides with the crossing of the dustlane and ring.

Zooming on the central arcminute, the lower right panels in Figure 5 show the uniform weight H I column density map superposed on the *R* band image (center right) and on the H α image (bottom right). The grayscale in the *R* band image is chosen to highlight the dustlane west of the nucleus. As seen in Figure 5, H α is detected only in the central regions, peaking at the nucleus and also in a ring of radius ~15". The ring of star formation is coincident with a shoulder on the *R*-band profile between 12 and 20". The H I in the inner regions avoids the nucleus but follows closely the H α within the limits of the VLA map resolution. Comparison of Figure 5 with Figure 1 of García-Burillo *et al.* (1998) reveals that the molecular gas peaks in the same overall ring, but just

TABLE 8
H I PROPERTIES

	U 6341	N 3626 U 6343	NPM1G +18.0219	N 3900 U 6786	N 4138 U 7139	N 4772 U 8021	N 5854 U 9726
S_{cont} (mJy)	< 0.3	9.9±0.1	< 0.3	1.2±0.1	32 ^b	3.4±0.2	...
$< DEF >^a$	...	+0.07	...	+0.02	+0.08	+0.20	1.62
$\int S_{HI} dV$ (Jy km s ⁻¹)	2.0±0.2	8.3±0.3	1.7±0.2	16.7±0.7	20.6 ^b	15.1±0.6	0.30 ^c
M_{HI} (10 ⁹ $M_\odot$)	0.27	1.1	0.23	2.9	1.2 ^b	0.9	0.04
M_{HI}/L_B ($M_\odot/L_\odot$)	...	0.06	...	0.2	0.2	0.1	.01
V_{21} (km s ⁻¹)	1645	1484	1466	1801	888 ^b	1040	1654 ^c
ΔV (km s ⁻¹) ^d	139	382	96	450	340 ^b	463	128 ^c
R_{HI}/R_{25}	...	2.6	...	1.8 ^e	2.5 ^b	1.9	...
Dynamical center (1950)	111722.8	111726.1	111807.3	114633.9	120658.3 ^b	125056.0	...
	+183202	+183748	+182938	+271757	+435749 ^b	+022621	...
Inclination (°) ^f	...	44–68	...	65	46–25 ^b	65–58	...
V_{max} (km s ⁻¹)	...	195	...	220	208 ^b	240	...

^aHI deficiency measure from Magri (1994).

^bFrom Jore *et al.* (1996).

^cBased on reanalysis of digital archive copy of Magri's spectrum.

^dWidth at 20% of peak of global profile, uncorrected for smoothing or turbulence.

^eFor main disk; including the southern extension, $R_{HI}/R_{25} \sim 4.5$

^fWhere a range is given, a flat rotation curve has been assumed, and the range indicates the variation from the inner to the outer disk. See Jore (1997).

where the H I and H α are not seen, and vice versa.

Figure 6 shows representations of the H I distribution and velocity field extracted from the low resolution (natural weight) column density map. In the three panels, the optical boundaries $D_{25} \times d_{25}$ are indicated by the ellipse, oriented along the optical major axis at $PA = 157^\circ$. The left panel shows the H I contours and their corresponding grayscale. The center panel shows the H I column density contours superposed on a grayscale representation of the R band image with the same scaling as in the center left panel of Figure 5. The H I is found in the central regions at $R < 60''$ and in a low column density ring of $N_H \sim 1\text{--}3 \times 10^{20}$ atoms cm^{-2} peaking at $R \sim 190''$. This outer ring is both exterior to the optical radius, significantly more elongated, and oriented with a different position angle, nearly north–south. The ratio of H I to optical size is large: $R_{HI}/R_{25} = 2.6$. The rightmost panel shows the H I velocity field derived from the natural weight data cube. The velocity field implies fairly regular rotation in the outer ring, with the north side of the galaxy is approaching and the south, receding. Many of the contours are jagged due to the patchy distribution of the gas. It is also apparent that the inner H I disk has a significantly different kinematic position angle than that of the outer disk or as measured from the overall gas distribution.

Two nearby galaxies were detected in the field of view of the H I map: UGC 6341, which is located 5.9 south of NGC 3626, and a faint object identified as NPM1G +18.0219 (Klemola *et al.* 1987) 12.9 (90 kpc) to the SE. In fact, the highest H I column density in the field is found coincident with the faint companion UGC 6341, an inclined low surface brightness Magellanic spiral. The H α image did not extend out this far but in the broadband images, two central knots are visible which makes fitting central ellipses impossible. There is no bulge evident. One of the knots appears to be the center of light (nucleus) and the other lies about $10''$ to the NW. We derive $m_R = 15.04 \pm 0.04$, with $\epsilon = 0.72$ and $PA = 153^\circ$. The global HI properties of these objects are summarized in Table 8 and the global H I profiles of all three objects are shown in Figure 4.

In the central regions, the uniform weight H I map is deep enough to show the column density contours, but the S/N is too low to trace the velocity field. Within the resolution, the dynamical center coincides with the nucleus, but the inner velocity field is clearly distorted. To the north, the velocity contours close

at $R \sim 12\text{--}25''$ but the velocity begins to rise again outside $45''$. This rise and subsequent fall appears to coincide with that seen in the optical emission line gas rotation curve, shown in Figure 7. To the south, even in the center, the velocity contours change continuously. Analysis of the velocity field was accomplished by following the common practice of fitting a tilted ring model, assuming circular rotation. In the center, the position angle of the H I is closely aligned with the optical axis, starting at 160° but swiftly increases to 195° at a radius of $65''$. The position angle then drops to 175° by a radius of $140''$ and changes little to the edge of the H I disk. Figure 7 shows position–velocity diagrams extracted from the natural weight data cube at position angles of 157° (upper left), along the optical major axis, and 178° (upper right), along the kinematic major axis of the outer H I ring. In the left panel, the optical stellar and emission line velocities derived from the long-slit spectra are superposed. It is clear that the H α and H I gas show similar rotation curves in the inner part, but that both counterrotate with respect to the stellar component. Moreover, it is also clear that the inner rotation curves of the ionized and neutral gas are both distinct and shifted in orientation from that traced by the outer H I ring. The rotation curve of the H I in the outer ring along $PA = 178^\circ$ continues to rise to its outermost point; the H I velocity gradient of the outer ring is minimized along $PA = 90^\circ$. A large velocity gradient is evident within the inner $\sim 20''$ in all cuts through the H I cube, due, at least in part, to beam smearing.

The H I velocity contours suggest that the inclination of the H I disk varies considerably. The best fit inclination given by the tilted ring model starts at 44° and then drops to 33° out to a radius of $80''$. However, this drop is likely an artifact of the disconnect between the inner H I ring and the outer H I ring; there is no H I in this region to constrain the velocity field. Towards the outside, the inclination appears to increase continually with radius until it reaches a maximum of 68° at the edge of the H I disk. That the outer ring is more highly inclined than the stellar disk is also suggested by the differences in ellipticity seen in the isophotes of optical light and H I column density. With this variation in inclination, the rotational velocity remains relatively constant at 195 km s^{-1} throughout the H I distribution.

Van Driel *et al.* (1989) obtained an H I synthesis map of this object with the WSRT. The significantly lower resolution ($13'' \times 41''$) and sensitivity of that

map did not allow discrimination of the structural details but there is rough correspondence to the features reported here. In fact, despite the poor resolution of their derived velocity field ($40'' \times 41''$), van Driel *et al.* noted the marked shift of position angle between the optical major axis and the morphological and kinematic position angles of the outer H I ring.

Ciri *et al.* (1995) obtained both absorption and emission line spectra at four position angles: 25, 68, 113 and 158° . They found that the stellar component entirely counter-rotates with respect to the stars. Along the $PA = 158^\circ$ axis, the stellar lines were traced to $\pm 30''$, while the ionized lines extended to $\pm 20''$. The Palomar spectra were obtained along $PA = 157^\circ$ (major axis) and $PA = 67^\circ$ (minor axis); as shown in the lower panels in Figure 7, the extracted rotation curves for both components are qualitatively similar to those shown in Figure 1 of Ciri *et al.* but extend further in the spatial dimension. The stellar curves rise with some deviation at about $8''$ to flatten at $R \sim 20''$ and beyond. The stellar curves can be traced in these spectra to $45''$ on either side of the major axis; this radius is only slightly more than $0.5 R_{25}$.

Emission line rotation curves can be traced to $\pm 20''$ in [OIII] and [SII] and to $\pm 30''$ in $H\alpha$ and [NII]. Although in the very center the $H\alpha$ is likely affected by absorption, there appears to be a hint that the $H\alpha$ emission retains its high velocity almost to the center, showing velocities of $\sim 150\text{--}180 \text{ km s}^{-1}$ within the inner $5''$. In fact, Ho *et al.* (1997a) classify NGC 3626 as a possible LINER, based on line ratios; the FWHM of the [NII] within their window along $PA = 170^\circ$ (Ho *et al.* 1995) is large, 413 km s^{-1} . The [NII], [SII] and [OIII] lines show steeply rising velocities along the major axis, but also evidence of line broadening or a secondary component is seen at symmetric points within $R \leq 6''$. In particular, the [NII] lines appear to double within the innermost region and have been fit by two separate Gaussian components, as indicated in Figure 7. When fit separately, the second component appears to match the velocity of the stellar component. In addition, both the $H\alpha$ and [NII] curves peak at symmetric radii at $R \sim 10\text{--}15''$ but then show a clear decline outside $R \sim 20''$. This decline is suggested but not evident in the lower sensitivity data of Ciri *et al.* (1995). As discussed above, these features are also reflected in the H I velocity field. The small deviation from constant velocity seen in the stellar values along the minor axis

is mostly likely due to a slight offset from the true dynamical center (i.e. pointing) or minor axis position angle. Along the minor axis, both the [NII] and [SII] spectra exhibit double-peaking within the inner $R \sim 6''$. Within the resolution limits, it is uncertain whether the line broadening is merely a LINER phenomenon, excess streaming, or a secondary gas component co-rotating with the stars. García-Burillo *et al.* (1993) likewise find suggestive evidence for a secondary gas component associated with the dominant stellar disk in the central regions.

In summary, the kinematic evidence for NGC 3626 suggests a large scale decoupling of the stars and gas. The H I distribution is found in two kinematically distinct and spatially separate rings that show a large difference in position angle and inclination. Both the morphological and kinematic position angles of the outer H I ring are shifted by $> 20^\circ$ with respect to starlight and the inner H I gas. Evidence of a gas component co-rotating with the stars is suggested, as hinted also in the CO map of García-Burillo *et al.* (1998).

3.2. NGC 3900

NGC 3900 is a ringed Sa galaxy with no nearby companions. The nearest known object of similar redshift, UGC 6791, lies to the SE at a projected separation of $17.1'$, 135 kpc at the adopted distance of 27 Mpc . NGC 3912, also considered part of a group LGG 252 by Garcia (1993) lies $34.8'$ (275 kpc) to the SW. There are several small galaxies nearby; two of them are included in the catalog of dwarfs found by Binggeli *et al.* (1990). None of these possible companions has a known redshift. For one, #44 of Binggeli *et al.*, we failed to detect any lines in a 600 sec exposure obtained with the Palomar 5m plus Double Spectrograph using the setup described in Section 2.2.

As evident in the B band images shown in Figure 8, NGC 3900 shows a small circular bulge with a prominent blue inner ring at a radius of $34''$. In the exterior, the light from the disk abruptly decreases at roughly half the optical radius. Flocculent spiral structure is clearly evident in a diffuse, faint disk that, as seen in the upper panel of Figure 8, appears to be more elongated than the higher surface brightness portions, as might be expected for a tightly wound spiral seen at moderate inclination. Only a hint of $H\alpha$ is evident in the narrow-band images with a single concentration near the center and a V-shaped structure in the north corresponding to blue enhancements ($B - V \sim$

0.6) in the northern section of the ring evident also in the color maps. A dustlane is seen on the southern side looping to the east and wrapping nearly halfway around the galaxy. The dustlane is easily visible on the northern and southern ends in the B band image, just interior to the inner ring and more easily distinguished in the east where it overlaps the inner ring in the color map images.

From examination of a HST WFPC2 F606W image, Carollo *et al.* (1997) classified N 3900 among the early spirals with a “classical spheroidal bulge”; the bulge component identified in the HST has a smaller effective radius than the limited resolution ground based images could distinguish from a central component. In fact, as evident in Figure 8, a $n = 2$ bulge fits quite well and gives a modest $B/D_R = 0.30$. Structure in the surface brightness profile is evident in all the broadband images where crossings of the ring and dustlane occur, and where the outer faint disk contributes. Use of a higher order bulge in our R band image yields a poorer fit with a higher $B/D_R = 0.71$ in which the bulge clearly overshoots the observed central light profile; the $n = 2$ bulge also provides a better fit to the outer ($R > 80''$) disk.

NGC 3900 was mapped previously with the WSRT by van Driel *et al.* (1989). In their map, most of the H I is found in a ring of radius $1'$, with some suggestion of the western extension. This ring is clearly evident in Figure 9 which shows different representations of the H I map and velocity field extracted from the natural weight VLA data cube. The H I indeed peaks in a ring at $R \sim 50''$, just outside the prominent blue ring, and in the region dominated by flocculent spiral structure. The lower sensitivity of the WSRT map completely missed the extended diffuse structure to the north and south, evident here in Figure 9. In addition to the higher density H I which is roughly coincident with the optical galaxy, diffuse H I can also be traced in patches to the north and south. To the south, an extension is visible, clumping about $5'$ (40 kpc) from the optical center. Furthermore, the H I column density distribution associated with the main body is clearly asymmetric, extending nearly twice as far to the west as to the east.

The velocity contours shown in the right panel of Figure 9 and the top left panel of Figure 10 clearly suggest that the southern gas cloud is simply an extension of the main H I distribution, the column density dropping below the sensitivity limit of the VLA map in between. In the inner parts, the contours in

the H I disk are very symmetric about the major axis and show regular rotation. The main H I disk extends only 2.9 (23 kpc) from the center; along the major axis, $R_{HI}/R_{25} = 1.8$. However, comparing the total extent of H I gas on the southern side of the galaxy to the stellar disk gives $R_{HI}/R_{25} = 4.5$; the H I in the south can be traced to $R \sim 50$ kpc. Along the minor axis, to the west, the H I extends about twice as far as to the east and deviates in velocity as would be expected if the H I disk were strongly warped in the outer parts.

The velocity field was modeled using tilted rings and including the southern low column density gas under the assumption that it lies in the same plane, at a constant inclination of 65° . In the best-fit model, the position angle varies from 178° to 186° in the main H I disk and then settles back to 176° at a radius of $200''$. With these parameters, the rotational velocity rises to ~ 200 km s $^{-1}$ in the inner $60''$ and is then relatively constant except between $240''$ and $300''$ where the parameters are poorly constrained because there is little gas in that region. In these models, the western extension is not well fit because of the asymmetry in the distribution along the minor axis.

Figure 10 shows the position-velocity diagrams for the H I distribution along both the optical major ($PA = 2^\circ$) and minor ($PA = 92^\circ$) axes; the velocities extracted from the long-slit spectra for stellar and gaseous components are superposed. The optical and H I rotation curves are similar, but the optical ones do not conclusively show the flattening seen in the H I data. There is also a suggestion of a spread in the velocity along $PA = 2^\circ$ to the north at $R \sim 1.5$, showing a blueshifted cloud with $\Delta V \sim 100$ km s $^{-1}$. In the upper right panel of Figure 10, the minor axis position-velocity diagram shows clear evidence of the H I ring and central hole. It also reveals a small change in velocity along the western H I extension, likely due to the proposed warp.

As illustrated in the lower panels of Figure 10, the optical spectra yield both gas emission and stellar absorption lines extending along the major axis to $R \sim \pm 55''$. The stellar Mg I b lines are relatively weak outside the bulge dominated region, at radii $> 10''$ to the north and $> 15''$ to the south. The stellar component to the N of the nucleus shows an asymmetric line of sight velocity distribution that could be either two separate components but is more likely simply due to true asymmetric deviation from a Gaussian distribution; this detail will be discussed elsewhere (Jore

et al. in preparation). The central H α is strongly affected by absorption within the inner 20'' to the north or 15'' to the south, but matches nicely the [NII] further out. In the inner rotation curves derived from the [NII], [OIII] and [SII] lines, a significant anomaly, indicating blueshifts of $\sim 160 \text{ km s}^{-1}$, is seen, peaking $\sim 5''$ to the north of center along the major axis. Beyond this radius, the velocities fall and resume the expected rise seen to the south at about 10''. There is a suggestion that the [N II] line is double peaked in the center. Ho *et al.* (1997a) classify NGC 3900 as a possible LINER based on line ratios extracted from a spectrum obtained along a PA of 178° (Ho *et al.* 1995). However, as can be seen in the lower panel of Figure 10, the [NII] lines exhibit a systematic sinusoidal wobble within $\pm 7''$ from the center along the minor axis. While this behavior might be expected from a small offset in position or position angle from the true kinematic center, the stellar velocities, extracted from the same spectra, remain flat. The color maps show a coincident feature — slightly bluer on the east and redder on the west of the nucleus at comparable radii.

While the main disk of NGC 3900 shows motions dominated by circular rotation in a quiescent pattern, anomalous emission line velocities imply non-circular motions in the central region. Interior to the optical edge, the H I is found in a ring or tightly wound spiral pattern with symmetric peaks to the north and south along the major axis in the region of flocculent optical spiral structure. At low column densities, $N_H \leq 2 \times 10^{20} \text{ atoms cm}^{-2}$, the H I distribution is huge. The outer H I can be traced to a radius more than 4 times the optical size in a diffuse and patchy distribution that appears kinematically connected to the main galaxy. Along the minor axis, the H I distribution is both warped and strongly asymmetric. Models of the evolutionary history of this galaxy must account for the kinematic quiescence of the patchy and asymmetric H I far beyond the optical edge.

3.3. NGC 4772

NGC 4772 is an outlying member of the Virgo cluster, probably associated with the subclump around NGC 4472. The adopted distance assumes this association and the derived subgroup distance of 16 Mpc from distance moduli given in Ferrarese *et al.* (2000). Its nearest companion, CGCG 015-036, a compact high surface brightness galaxy with a systemic velocity of $V_\odot = 844 \text{ km s}^{-1}$, is found at 18'.5, 85 kpc

at the adopted distance of 16 Mpc to the E-SE. The pair NGC 4809/4810 at similar velocity lies 35' (160 kpc) to the NE.

Dubbed the “eye galaxy,” a descriptive term adequately describing its appearance as shown in Figure 11, NGC 4772’s center is dominated by a slightly flattened bulge surrounded further out by a segmented ring. The western and eastern sides of the ring form continuous structures; however, the northern and southern ends are broken and patchy. The inner galaxy shows a clear dustlane surrounding the bright bulge and oriented parallel to the major axis. The bright, red circular bulge, of $B - V \sim 1.0$, is clearly separable within the inner 5'', cutoff on the north and east by the dustlane. Structure (filaments or braids) are seen within the dustlane, which must trace spiral structure, not simply a ring. The bulge-disk decomposition yields a moderate $B/D_R = 0.44$.

The H α shows a central peak, coincident with the nucleus, surrounded by an elongated ($b/a \sim 0.3$) H α ring which itself is crisscrossed but entwined in the dustlane and which coincides with the H I column density peaks. As evident in the upper panel in Figure 11, there are two pronounced drops in surface brightness, one at about 65'' and a second at about 150''. This latter coincides with the outer edge of the inner H I ring, just about where there are suggestions of the start of spiral arms. At that radius, the isophotes, $\mu_B = 25.9$, $\mu_R = 24.0 \text{ mag arcsec}^{-2}$, become rounder, changing from $\epsilon \sim 0.47$ to $\epsilon \sim 0.29$, similar to what is seen in the H I distribution. At the faintest isophotes, this rounder low surface brightness distribution can be traced to more than 200'', twice R_{25} , and roughly to the outermost H I contours. The outer low surface brightness disk is blue, $B - V \sim 0.67$ versus $B - V \sim 0.75 - 0.85$ further in, but contributes only a small amount of light (0.19 mag in B) to the total luminosity. The faint optical light is visible on the DSS image but makes the galaxy rounder than published axial ratios suggest. It is reminiscent of a similar feature detected by Buta *et al.* (1995) in deep photometry of the counter-rotating Sab galaxy NGC 7217.

As evident in Figure 12, the H I gas is distributed in two distinct rings surrounding a central hole. Coincident with the H α ring evident in Figure 11, the inner high column density H I ring peaks at $R \sim 60''$, with a position angle of 144° , similar to that of the bright optical disk and with comparable eccentricity. Surrounding the central hole, the full ring is charac-

terized by moderate column densities, $N_H > 7 \times 10^{20}$ atoms cm^{-2} . Clearly separated from this inner ring, the rounder outer H I ring can be traced to $\sim 200''$ in radius oriented along a position angle of 150° and, as seen in the lower right panel of Figure 12, extends over an area similar to, but of different eccentricity, with the faint blue stellar light. The outer ring is patchy, but, in the north, can be traced over 90° in azimuth at $N_H \sim 2 \times 10^{20}$ atoms cm^{-2} , with localized patches of 3×10^{20} atoms cm^{-2} . The velocity field shows a similar discontinuity. Near the edge of the inner ring and in the outer ring, along the major axis, the isoveLOCITY contours bend, suggesting a warp in the outer regions. Figure 13 shows the position-velocity diagrams along the optical major (PA= 147°) and minor (PA= 57°) axes; the velocities derived from the optical spectra in the same regions are shown. The velocity rises linearly in the inner regions and then flattens out at a radius of $90''$. Within the outer ring, the H I velocities show a significant turn-over at large radii in the SE. Along the minor axis, the contours appear regular and symmetric surrounding the central hole, but the gradient of the outer H I velocities reinforces the notion that the outer gas disk is warped.

As in the cases of NGC 3626 and NGC 3900, the fitting of tilted ring models is complicated by the discontinuity in the H I distribution between the inner and outer rings. The inner region shows a rotational velocity smoothly rising to 240 km s^{-1} at a radius of $100''$. If a flat rotation curve, set by the fit to the inner disk, is adopted, the position angle starts at 143° and rises steadily to 160° near a radius of $140''$ remaining roughly constant at $\sim 158^\circ$ further out. However, just at the edge of the inner H I ring, the inclination becomes unconstrained and discontinuous. A second fit was performed to smooth over the discontinuity in inclination by fixing the inclination for the transition rings to allow a smooth variation in the inclination. In that case, the inner rotational velocity rises much in the same way as in the previous model, but in the outer ring, it rises by 60 km s^{-1} . To maintain a flat rotation curve, the inclination must drop outside $R > 110''$ from about 65° to about 58° , as suggested also by the rounding of the aspect ratio of the outer column density contours. In terms of residuals, both of these models fit the observed velocity field equally well. Further details of the resultant mass modelling will be discussed elsewhere.

Figure 13 also shows the stellar and ionized gas velocities extracted from the long slit spectra. The

stellar lines can be traced from about $60''$ on the SE to about half as far on the NW. The ionized gas distribution is extremely patchy. [NII], [OIII] and [SII] emission is detected within the central $R \leq \pm 6''$, and on the southeast side along the major axis in clumps at $R \sim 25''$ and $50''$. No ionized emission is detected to the NW. The $\text{H}\alpha$ line is depressed in the center due to absorption as far out as $15''$ but the outer $\text{H}\alpha$ emission coincides nicely with that of the other ionized species. As also evident in the top panels of Figure 13, the optical rotation curves do not extend sufficiently far to sample the rotation curve peak.

Ho *et al.* (1997ab) classify NGC 4772 as a Seyfert Sy1.9, based on line ratios and the appearance of broadened $\text{H}\alpha$ seen in their spectrum obtained along PA = 5° . They also remark on the flat or double-peaked nature of the narrow lines. The $\text{H}\alpha$ line broadening is also evident in our spectrum but the other species, particularly [OIII], show a systematic rise to $\sim 80 \text{ km s}^{-1}$ at $R \sim \pm 6''$, more steeply and of opposite sign than the stellar rise in the same region. It is important to note that, unlike Ho *et al.*, who chose to orient the slit close to the parallactic angle, our setup placed the slit along the optical axis at P.A. = 147° . Furthermore, as evident in Figure 13, the minor axis [NII], [OIII] and [SII] velocities rise rapidly in the inner $5''$ up to a velocity of 100 km s^{-1} and then decrease out to $15''$ in radius. In this case, the [OIII] provides the best tracer, showing a velocity peak along the minor axis at $4-5''$ followed by a decline back to the systemic by the time it disappears at about $15''$. At the same time, the minor axis stellar velocities are very flat (verifying that the slit was correctly aligned along the minor axis of the *stellar* component). The kinematic signature suggests the presence of a misaligned disk or bar. The only unique feature seen in the color maps is a flattening of the $B-R$ in the central region along the minor axis and a hint of smaller plateaus in all the color maps along the major axis. These two color features occur on comparable scales as the central rotation curve peculiarities.

NGC 4772 may represent the end-state of a prograde merger in which the transfer of angular momentum leads to an outward spread of the disk (Quinn *et al.* 1993). The steep rise of the ionized gas velocities along the minor axis and the apparent decoupling on gas emission along the major axis suggests the presence of a misaligned embedded disk or bar. Reminiscent of NGC 3626, the H I is contained in two separate

but concentric rings of differing position angle and eccentricity. The inner ring coincides with the sites of current star formation in the optically bright inner disk, while the other rounder one, wholly outside R_{25} but coincident with a faint blue stellar component.

3.4. NGC 5854

NGC 5854 is a gas-poor Sa with no evidence of current star formation. Relatively isolated, it is likely an outlying member of the NGC 5846 group (Haynes & Giovanelli 1991; Garcia 1993; Zabludoff & Mulchaey 1998). Its nearest neighbor is CGCG 021-009 at $33'.1$, 230 kpc at the assumed distance of 24 Mpc. An illustrative summary of results is presented in Figure 14.

The B band image shows an inclined galaxy with a large elongated bulge and little apparent spiral structure. Some hint of arms is seen in the two small stumps on the NE and SW edges of the bulge. Unfortunately, the B band image was taken under non-photometric conditions so that the color information is limited to $V - R$. In the $V - R$ image, little structure is seen, although the galaxy is clearly redder than the others. It was previously imaged by Balcells & Peletier (1994) who find $B/D_R = 0.46$ using a 1-D decomposition method and find similar flatness of the color profile. As discussed below, no $H\alpha$ is seen in the long slit spectra, but $H\alpha$ absorption is seen across the disk and can be traced almost as far as the Mg I b lines. The galaxy is thus not currently forming stars, but its significant population of A stars suggests a post-starburst state. Of the galaxies discussed here, it has the highest $B/D_R \sim 0.6$ for $n = 2$, but it is clear that the $n = 4$ fit, illustrated in Figure 14, gives a more appropriate $B/D_R \sim 0.8$. No $H\alpha$ image was obtained.

As reported by Magri (1994), the H I mass of NGC 5854 is very low ($\sim 3 \times 10^7 M_\odot$) and hence it was not feasible to map its H I with the VLA. The optical spectra are dominated by stellar absorption lines which can be traced, by both Mg I b and $H\alpha$, out to $\pm 45''$, further than, but qualitatively similar to the rotation curve displayed in Simien & Prugniel (1997). Those authors note $V_{max} = 128 \pm 8 \text{ km s}^{-1}$ for the stellar component in their spectrum traced to $\sim 27''.5$. We find a comparable value at that radius, but a larger $V_{max} = 167 \pm 10 \text{ km s}^{-1}$ at $R \sim 40''$. The Mg I b rotation velocities rise more shallowly than the $H\alpha$ lines do, possibly because of extinction. The [NII] and [OIII] emission reveals a distinct counter-rotating

gas component traceable from the center out to $6''$ on the NE side and $8''$ on the SW side. The maximum rotational velocity of this [NII] feature is 64 km s^{-1} at $\sim 4''$ NE of the optical center of light, the [OIII] appears in a ring coextensive with the [NII] gas but the velocities are higher, reaching $\sim 100 \text{ km s}^{-1}$. The stellar velocity profiles along the minor axis show a slight trend that most likely arises because of a small pointing offset. The weak [NII] shows some dispersion but is essentially flat, within the errors; no other ionized emission is seen.

Magri (1994) detected a weak and narrow H I signal from NGC 5854 at Arecibo after more than 2 hours of on-source integration. A zoomed reproduction of Magri's spectrum extracted from our digital archive is shown in Figure 14. The systemic velocity of the H I is 1654 km s^{-1} which matches the value of $1663 \pm 10 \text{ km s}^{-1}$ given by Simien & Prugniel (1997). We have remeasured the H I line width in the digital spectrum and corrected it for smoothing, etc. following Haynes *et al.* (1999b) to obtain a value, uncorrected for inclination and turbulence, of $W_{21} \sim 95 \text{ km s}^{-1}$. This value is too small compared to the full rotation width of the galaxy, but is comparable to the velocity spread seen in the [NII] in the inner disk. A simple interpretation of the narrowness of the H I profile is that the H I and [NII] both arise in a small central disk of $R < 8''$ that counter-rotates relative to the stars. At the same time, the primary (stellar) disk is seen in a post-starburst phase, showing strong Balmer absorption throughout. The current gas-poor state of NGC 5854 may thus be result of the accretion of a relatively dense satellite that has heated the disk and smoothed the spiral structure, leaving only the counter-rotating gas core as evidence of its occurrence.

3.5. Another Look at NGC 4138

As mentioned previously, the first object mapped in H I with the VLA as part of this study was NGC 4138. A discussion of the optical and H I spectral observations is presented in Jore *et al.* (1996); the gas counterrotates with respect to the primary stellar component, but a secondary stellar population, kinematically coupled to the gas, is visible. The amplitude of the crosscorrelation function of the stellar lines coincides spatially with an $H\alpha$ ring, previously noticed by Pogge & Eskridge (1987), at a radius of $\sim 20''$. The two stellar components are characterized by distinct velocity dispersions, implying separate ori-

gins. As part of the current study, we added B , V , R and $H\alpha$ images to the dataset for this morphologically normal but kinematically unusual object.

As evident in the top left panel of Figure 15, NGC 4138 is a small bulged, smooth armed Sa galaxy with a prominent dustlane in the SE quadrant. A foreground star is located at a radial distance of $4''$ from the center along the major axis. As seen in the middle panels of Figure 15, the inner H I forms a partial ring, coincident with the $H\alpha$ ring, on the north and east, surrounding a central hole. In $B - V$, the galaxy shows a distinctly red bulge ($B - V \sim 1.1$) and a blue ($B - V \sim 0.6$) ring located in the approximate position of the $H\alpha$ ring. The blue ring is particularly strong on the northern and southern portions of the galaxy with breaks in the ring on the east and west. The eastern break is due, at least in part, to the dustlane. The $V - R$ image shows similar structure. As shown in the upper right panel of Figure 15, the bulge–disk decomposition gives a small value: $B/D_R = 0.17$.

There are several outstanding features in the color profiles which correlate well with features in its rotation curve as presented in Jore *et al.* (1996). At $\pm 13''$, where the secondary component of stars begins, until $\pm 35''$ when the component is no longer detectable in the long slit spectrum, there is an obvious reddening in the major axis profiles. There is an abrupt transition moving from the center into the ring on the southeast side and an abrupt transition out of the ring on the northwest side. There are no apparent variations in the color profiles along the minor axis aside from a peak redder color in the center.

The relationship of the H I gas reported previously by Jore *et al.* to the ring and the dustlane is demonstrated here in more detail in Figure 15. Indeed, the relationship of the dustlane and $H\alpha$ ring to the H I distribution seen in NGC 4138 is very similar to that seen in NGC 3626 (Figure 15 versus Figure 5), the other case of large-scale counter-rotation. Likewise, the H I extent is very large, $R_{HI}/R_{25} = 2.5$ and, as discussed in Jore *et al.*, the velocity contours along the minor axis suggest a strong warp beyond R_{25} . In contrast to the H I distributions seen in NGC 3626 and NGC 4772, however, the outer HI does not form a distinct and coherent ring, but rather is extremely patchy beyond the optical edge.

NGC 4138 remains the sole example in our Sa sample of a galaxy with an extended counter-rotating stellar component. Nonetheless, it bears similarities

to the other objects in terms of its morphological “boring-ness”, relative isolation, large, low surface density H I extent, and overall regularity of its velocity field.

4. Discussion

In this paper, we have presented a combined imaging and spectroscopic dataset for four Sa galaxies; images are also included for a fifth, NGC 4138, discussed previously by Jore *et al.* (1996). All of the galaxies show evidence of departure from kinematic normalcy ranging from large-scale counter-rotation to decoupled central gas and/or stellar components. Despite the difference of detail, several unifying themes seem critical: (1) All of the objects are relatively isolated, morphologically normal, unbarred Sa galaxies. (2) Rings are prominent features both in the optical broadband light and in the distributions of ionized and neutral gas. The apparent rings may also result from tightly wound spiral arms viewed at moderate inclination. (3) The kinematically decoupled gas components appear to be associated with sites of current or at least recent star formation. In all cases where $H\alpha$ emission is detectable in the region of interest, it coincides well with the location of kinematically decoupled components. When there is no $H\alpha$ emission in these regions, the decoupled components often show some form of a color change, often towards the red. (4) The gas-rich galaxies (all except NGC 5854) contain moderate H I masses, but, because the gas is spread over a large area $R_{HI}/R_{25} \geq 2$, the globally averaged H I surface densities $\langle \sigma_{HI} \rangle = M_{HI}/\pi R_{HI}^2$ are very low, typically of order $0.5\text{--}1.0 M_{\odot} \text{ pc}^{-2}$. In NGC 3626 and NGC 4772, the H I is found in two concentric but distinct rings. In NGC 3900, the gas exterior to the optical edge is patchy, but can be traced 50 kpc to the south, giving $R_H/R_{25} \sim 4.5$. Along the minor axis, the exterior H I is strongly asymmetric. (5) In all four H I maps, including that of NGC 4138, the velocity field is dominated by circular rotation, but significant departures from motion in a quiescent disk are evident. In all, the outer H I velocity contours suggest significant warping of the outer disk, roughly beginning at the optical edge and following Briggs’ (1990) rule for warps, as commonly seen in other galaxies. The similarities found in the five Sa galaxies are consistent with the scenario that the kinematically distinct gas components arise from slow, minor mergers of gas-rich satellites with an already-formed disk galaxy. Differences in their appearance

thus arise from differences in the circumstances of the merger events. Retrograde primordial gas infall, as discussed by Thakar *et al.* (1997) is also offered as a possible solution to produce counter-rotating components, although seemingly more difficult to justify in the growing number of known cases.

Though most numerical work on the merger process has focussed on equal mass progenitors, recent studies have begun to address the minor merger phenomenon (Quinn *et al.* 1993; Hernquist & Mihos 1995; Walker *et al.* 1996; Thakar *et al.* 1997; Bekki 1998; Thakar & Ryden 1998; Taniguchi 1999). Although much of this work is preliminary, requiring further refinement of the range of galaxy characteristics and orbital parameters and the influence of gas dissipation, triaxiality and dark matter content, we can nonetheless examine the main characteristics of the five galaxies under the assumption that we may be seeing the results of varying acquisition scenarios.

The extensive H I distributions seen in four of the five galaxies discussed here may result from the slow accretion of a gas-rich satellite that is tidally stripped before the merger occurs. Accretion of a low density gas-rich satellite would minimize the disk heating problem and might favor the formation of extended H I disks or rings, but perhaps not inner rings. The formation of rings is seen in a variety of infalling gas models, both due to orbit crowding and to true self-gravitating rings. Also, Quinn *et al.* (1993) suggest that transport of angular momentum outwards will tend to cause the disk to spread in radius. The newly-acquired gas may reach sufficient densities, if clumpy, to form a new generation of stars. This possibility is suggested by the faint, blue, low-surface brightness outer disk seen in NGC 4772 and a similar excess of light found around the counter-rotating Sab galaxy NGC 7217 (Buta *et al.* 1995).

Clearly the timescales of the merger events responsible for the individual characteristics noted here are likely to vary greatly. Relatively gas-poor and large-bulged, NGC 5854 may represent the result of the accretion of a more massive satellite travelling in a prograde orbit (Walker *et al.* 1996). Merger events might lead to bulge build-up and even induce large-scale star formation, leaving the galaxy in the post-starburst state in which NGC 5854 is found. Furthermore, one of the primary arguments made against the tidal forcing of warps is that the modes would damp out after a few galactic rotations. Perhaps in the cases that suggest strong warping of the H I, the

minor mergers may be relatively recent events. Over time, the warps may damp, or be reexcited by additional interactions. If there are no impediments to star formation in a counter-rotating gas component, NGC 4138 with its two counter-rotating extended stellar disks may represent the future evolutionary state of NGC 3626.

The similarities among the results for these five galaxies lead to several questions and conundrums. (1) Is it key that the sample is composed of Sa galaxies? The accretion of moderate amounts of gas may also sweep up the gas contained within the primary disk itself, ultimately leading to a smoothing of the spiral structure. Do minor mergers lead to bulge build-up and disk heating, driving the post-merger morphology towards earlier spiral types? (2) Damping the process of disk heating is a challenge for models of mergers, even minor ones, if the accreted satellite is sufficiently massive or dense. The resultant disk heating will be relatively lessened if the satellite is predominantly gaseous. What are the conditions necessary to avoid over-heating the disk and disturbing the overall velocity field and spiral structure? (3) Is the fact that these galaxies are all unbarred important? Loss of angular momentum may drive the material inward toward the central regions where it accumulates in a nuclear disk as is often discussed for the kinematically decoupled cores seen in ellipticals. In the early stages of galaxy evolution, the formation of bars seems inevitable, but if the central concentration of mass is sufficient, the development of an inner Lindblad resonance will inhibit the central flow of gas (Sellwood & Moore 1999). Thus, since the progenitor primary was already a well-developed object, infalling material may be stopped at the inner Lindblad resonance, allowing the build-up of gas in a ring with subsequent star formation. Does this process produce the inner H α and H I rings seen here? Jungwiert & Palous (1996) propose that unbarred ringed galaxies must possess weak central oval distortions, while the two-stream instability picture of Lovelace *et al.* (1997) for galaxies with counter-rotating components suggests the generation of strong m=1 spiral waves. Are the inner kinematic peculiarities evidence of such oval distortions or m=1 streaming? (4) Although these Sa galaxies are relatively isolated, they are all members of loose groups. In fact, Sa galaxies are found typically in denser environments than their later type counterparts. Is the relative morphological segregation seen across the spiral sequence related to

the likelihood of minor mergers and their increased probability in environments characterized by moderate density and low velocity dispersion?

While examination of the rate of strongly disturbed galaxies may provide a reliable estimate of the major merger rate (e.g. Keel & Wu, 1995), minor mergers may be morphologically hidden, revealed only by the presence of their resultant kinematic peculiarities. Because of the ambiguities involved in disentangling the kinematic clues, it is not possible to trace with any degree of precision the evolutionary history of the multiple stellar and gaseous components evident in these galaxies. Food for thought, however, is the possibility that, over the course of its lifetime, a disk galaxy might undergo multiple minor accretion events with kinematic as well as morphological consequences. In particular, more complicated histories involving multiple minor mergers are likely to produce heterogeneous properties, particularly those related to properties of “form”. We suggest that the heterogeneity of the Sa class in particular is the outcome of such complicated life histories, the memory of which is signaled only by kinematic clues such as those identified here.

We thank Giovanni Moriondo for providing his 2-D bulge-decomposition code FITMAP and Giovanni Moriondo, David E. Hogg and Morton S. Roberts for numerous discussions. This work has been supported by NSF grants AST-9023450 and AST-9528860 to MPH. Portions of this research have been used in partial fulfillment of the Ph.D. (KPJ) and M.Eng. (EAB) degrees at Cornell University. BMM was supported by NSF-REU grant AST-9619531. This study has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, Caltech, under contract with the National Aeronautics and Space Administration and the Digital Sky Survey which was produced at the Space Telescope Science Institute under US Government grant NAG W-2166.

REFERENCES

- Balcells, M. and Peletier, R. F. 1994, *AJ*, 107, 135
- Bekki, K. 1998 *A&A*, 334, 814
- Bertola, F., Cinzano, P., Corsini, E. M., Pizzella, A., Persic, M. and Salucci, P. 1996, *ApJ*, 458, L67
- Binggeli, B., Tarenghi, M. and Sandage, A. 1990, *A&A*, 228, 42
- Briggs, F. H. 1990, *ApJ*, 352, 15
- Buta, R., van Driel, W., Braine, J., Combes, F., Wakamatsu, K., Sofue, Y., and Tomita, A. 1995, *ApJ*, 450, 593
- Carollo, C. M., Stiavelli, M., de Zeeuw and Mack, J. 1997, *AJ*, 114, 2336
- Ciri, R., Bettoni, D. & Galletta, G. 1995, *Nature*, 375, 661
- Corsini, E. M., Pizzella, A., Funes S. J., J. G. Vega Beltran, J. C., and Bertola, F. 1998, *A&A*, 337, 80
- de Jong, R.S. 1996, *A&AS*, 118, 557
- de Vaucouleurs, G., de Vaucouleurs, A., Corwin, H. G., Buta, R. J., Paturel, G., and Fouque, P. 1991. *Third Reference Catalogue of Bright Galaxies*. New York: Springer (RC3)
- Ferrarese, L., Ford, H. C., Huchra, J., Kennicutt, R.C. Jr., Mould, J. R., Sakai, S. *et al.* 2000, *ApJS*, (in press)
- Garcia, A. M. 1993, *A&A*, 100, 47
- García-Burillo, S., Sempere, M. J. & Bettoni, D. 1998, *ApJ*, 502, 235
- Haynes, M. P. & Giovanelli, R. 1984, *AJ*, 89, 758
- Haynes, M. P. & Giovanelli, R. 1991, *AJ*, 102, 841
- Haynes, M. P., Giovanelli, R., Salzer, J. J., Wegner, G., Freudling, W., da Costa, L. N., Herter, T. and Vogt, N. P. 1999a, *AJ*, 117, 1668
- Haynes, M.P., Giovanelli, R., Chamaraux, P., Freudling, W., da Costa, L.N., Salzer, J.J., and Wegner, G., 1999b, *AJ*, 117, 2039
- Hernquist, L., & Mihos, J. C. 1995, *ApJ*, 448, 41
- Ho, L. C., Filippenko, A. V. & Sargent, W. L. W. 1995, *ApJS*, 98, 477
- Ho, L. C., Filippenko, A. V. & Sargent, W. L. W. 1997a, *ApJS*, 112, 315
- Ho, L. C., Filippenko, A. V., Sargent, W. L. W. & Peng, C.Y. 1997b, *ApJS*, 112, 391

- Jore, K. P. 1997, Ph.D. thesis, Cornell University
- Jore, K. P., Broeils, A. H., & Haynes, M. P. 1996, AJ, 112, 438
- Jore, K. P., Haynes, M. P., Barrett, E. A. & Broeils, A. H. 2000, in preparation.
- Jungwiert, B. & Palous, J. 1996, A&A, 311, 397
- Keel, W. C. & Wu, W. 1995, AJ, 110, 129
- Klemola, A. R., Jones, B. F. & Hanson, R. B. 1987, AJ, 94, 501
- Laine, S. & Heller, C. H. 1999, MNRAS, 308, 557
- Landolt, A. U. 1992, AJ, 102, 340
- Lovelace, R. V. E., Jore, K. P. and Haynes, M. P. 1996, ApJ, 475, 83
- Magri, C. 1994, AJ, 108, 896
- Merrifield, M. R. & Kuijken, K. 1994, ApJ, 432, 575
- Mihos, J. C. & Hernquist, L. 1994, ApJ, 437, 611
- Mihos, J. C., Walker, I. R., Hernquist, L., Mendes de Oliveira, C. and Bolte, M. 1995, ApJ, 447, L87
- Moriondo, G., Giovanardi, C. & Hunt, L. K. 1998, A&A, 130, 81
- Pogge, R. W. & Eskridge, P. B. 1987, AJ, 93, 291
- Quinn, P. J., Hernquist, L. and Fullagar, D. 1993, ApJ, 403, 74
- Rieke, G. H. & Lebofsky, M. J. 1985, ApJ, 288, 618
- Roberts, M. S. & Haynes, M. P. 1994, ARA&A, 32, 115
- Sandage, A. & Tammann, G.A. 1987, *Revised Shapley Ames Catalogue of Bright Galaxies*, Carnegie Inst. of Washington Publ. 635, (RSA)
- Sellwood, J. A., & Moore, E. M. 1999, ApJ, 510, 125
- Simien, F. & Prugniel, P. 1997, A&A, 126, 158
- Taniguchi, Y. 1999, ApJ, 524, 65
- Thakar, A. R., Ryden, B. S., Jore, K. P., & Broeils, A. H. 1997, ApJ, 479, 702
- Thakar, A. R. & Ryden, B. S. 1998, ApJ, 506, 93
- van der Hulst, J.M., Terlouw, J.P., Begeman, K.G., Zwitter, W. & Roelfsema, P.R. 1992, *Astronomical Data Analysis Software and Systems I*, A.S.P. Conference Series 25, 131
- van Driel, W., Balkowski, C. and van Woerden, H. 1989, A&A, 218, 49
- Walker, I. R., Mihos, J.C., and Hernquist, L. 1996, ApJ, 460, 121
- Zabludoff, A. & Mulchaey, J. 1998, AJ, 496, 39
- Zaritsky, D. & Rix, H.-W. 1997, ApJ, 477, 118
- Zaritsky, D., Smith, R., Frenk, C. and White, S. D. M. 1997, ApJ, 478, 397

This 2-column preprint was prepared with the AAS L^AT_EX macros v4.0.

Fig. 1.— Cleaned continuum-subtracted H I line channel maps of NGC 3626 derived from the natural weight (low resolution) data cube. Contour levels range from 0.46 to 4.6 mJy beam⁻¹ in increments of 0.46 mJy beam⁻¹. The beam size is 16".8 x 16".2.

Fig. 2.— Cleaned continuum-subtracted H I line channel maps of NGC 3900 derived from the natural weight (low resolution) data cube. Contour levels range from 0.42 to 6.42 mJy beam⁻¹ in increments of 1.0 mJy beam⁻¹. The beam size is 22".5 x 20".0.

Fig. 3.— Cleaned continuum-subtracted H I line channel maps of NGC 4772 derived from the natural weight (low resolution) data cube. Contour levels range from 0.62 to 8.62 mJy beam⁻¹ in increments of 1.0 mJy beam⁻¹. The beam size is 31".7 x 23".1.

Fig. 4.— Global profiles derived from VLA mapping for the target galaxies and neighboring companions. H I parameters are given in Table 8.

Fig. 5.— Imaging results for NGC 3626. Top left panel: *R* band image with the scaling set to showing the maximum extent of starlight. Middle left panel: Expanded *R* band view illustrating the faint ring and spiral structure. The ellipse has the dimensions of $D_{25} \times d_{25}$ and is oriented along the optical major axis. Lower left panel: The inner 1' shown to emphasize the dustlane west of the nucleus. Upper right panel: Results of the two-dimensional bulge-disk decomposition for a $n = 2$ bulge. The solid line traces the light distribution while the disk and bulge are shown as long-dashed and short-dashed lines respectively. Middle right panel: The high resolution H I column density map superposed on a greyscale representation of the central *R* band image. The first H I contour is at 1.0×10^{20} atoms cm⁻². The next six contours range from 4.0 to 11.5×10^{20} atoms cm⁻² in steps of 1.5×10^{20} atoms cm⁻² and the remainder range from 13.5 to 18.5×10^{20} atoms cm⁻² in steps of 1.0×10^{20} atoms cm⁻². Lower right panel: Identical contours of the high resolution H I column density map superposed on the H α image.

Fig. 6.— H I mapping results obtained from the low resolution cube for NGC 3626 and UGC 6341. Left panel: H I column density contours and grayscale ranging from 1.0 (2σ) to 17.5×10^{20} atoms cm⁻² in steps of 1.5×10^{20} atoms cm⁻². Middle panel: The same H I column density map overlayed on the *R* band image. The optical greyscale has the same shading as in the center left panel of Figure 5. Right panel: The H I velocity field map. Velocity contours range from 1275 (light) to 1675 (dark) km s⁻¹ in increments of 25 km s⁻¹. In all three panels, the ellipse outlines the optical galaxy as given by $D_{25} \times d_{25}$ oriented along the optical major axis, PA = 157°.

Fig. 7.— Position-velocity cuts obtained from the low resolution H I cube and the optical spectra for NGC 3626. Upper panels: H I position-velocity diagrams along four slices through the low resolution data cube. The top left panel shows the slice along the optical major axis, PA = 157°, along which the optical spectra were obtained. In addition to the H I contours, the velocities of the optical stellar absorption line (open squares) and emission line (filled circles) are also shown. The secondary gas component is represented by open circles. The top right panel shows the slice at PA = 178°, the major axis of the outer H I distribution. The center left panel shows the slice along the optical minor axis, PA = 67°; the velocities of the stellar and ionized gas components obtained from the long slit spectra are also shown. The center right panel shows the slice along a position angle of 90°, the minor axis of the outer H I ring where the velocity gradient is minimized. The lower panels show the optical rotation curves derived for NGC 3626 along the major axis (PA = 157°) and along the minor axis (PA = 67°). Separate symbols show the rotation curves derived from the stellar absorption and gas components. Error bars are shown. In the innermost regions, the [NII] lines appear double.

Fig. 8.— Results of optical imaging for NGC 3900. Upper panel: B band image with scale set to emphasize outer structure. The ellipse has the dimensions of $D_{25} \times d_{25}$ and is oriented along the optical major axis, $PA = 2^\circ$. Middle panel: Low contrast B band grayscale showing spiral structure and dustlanes in the inner galaxy. The spatial scale is zoomed by a factor of two relative to the upper panel. Lower panel: Results of the two-dimensional bulge-disk decomposition for a $n = 2$ bulge. The solid line traces the light distribution while the disk and bulge are shown as long-dashed and short-dashed lines respectively.

Fig. 9.— H I mapping results obtained from the low resolution cube for NGC 3900. Left panel: H I column density contours ranging from 0.3 (2σ) to 12.3×10^{20} atoms cm^{-2} in steps of 2.0×10^{20} atoms cm^{-2} . Middle panel: Similar H I column density map, with a stepsize of 1.0×10^{20} atoms cm^{-2} overlayed on the B band image. Right panel: The H I velocity field map showing velocity contours ranging from 1575 (light) to 2050 (dark) km s^{-1} in increments of 25 km s^{-1} . In all three panels, the ellipse outlines the optical galaxy as given by $D_{25} \times d_{25}$ oriented along the optical major axis $PA = 2^\circ$.

Fig. 10.— Position-velocity cuts obtained from the low resolution H I cube and the optical spectra for NGC 3900. Top panels: H I position-velocity diagrams along two slices through the low resolution data cube. The top left panel shows the slice along the optical major axis, $PA = 2^\circ$, along which the optical spectra were obtained. In addition to the H I contours, the rotation curves traced by the optical stellar absorption lines (open squares) and the gas emission lines (filled circles) are also shown. The top right panel shows the slice at $PA = 92^\circ$, along the minor axis. The lower panels show the optical rotation curves derived for NGC 3900 along the major ($PA = 2^\circ$) and minor ($PA = 92^\circ$) axes. Separate symbols show the rotation curves derived from the stellar absorption and gas components. Error bars are shown.

Fig. 11.— Imaging results for NGC 4772. Top left panel: B band image scaled to show the faint outer disk. Middle left panel: B -band image of the same region but with the grayscale set to emphasize the inner regions, showing that the dustlane cuts across the bulge in the NE. Lower left panel: $H\alpha$ image. The ellipse superposed on the three images has the dimensions of $D_{25} \times d_{25}$ and is oriented along the optical major axis, $PA = 147^\circ$. Upper right panel: Results of the two-dimensional bulge-disk decomposition for a $n = 2$ bulge. The solid line traces the light distribution while the disk and bulge are shown as long-dashed and short-dashed lines respectively. Middle right panel: The high resolution H I column density map superposed on a grayscale representation of the B band image in the inner regions. Contours extend from 0.6×10^{20} to 14×10^{20} atoms cm^{-2} in steps of 4.0×10^{20} atoms cm^{-2} . Lower right panel: Identical contours of the high resolution H I column density map superposed on the $H\alpha$ image.

Fig. 12.— Results for the large-scale H I mapping of NGC 4772. Upper left panel: H I column density contours superposed on the B band image. Contours are shown at levels of $0.5, 1.5, 2.5, 4.0, 6.0, 8.0, 10.0$ and 12.0 and 14.0×10^{20} atoms cm^{-2} . Upper right panel: Grayscale and contours of the H I column density as shown in the left panel. Lower left panel: The outermost H I column density contour at 5×10^{19} atoms cm^{-2} superposed on a grayscale representation of the B band image scaled to enhance the faint outer disk. Lower right panel: The velocity field of NGC 4772, shown as contours and grayscale extending from 770 km s^{-1} (NW) to 1250 km s^{-1} (SE) in steps of 30 km s^{-1} . In all panels, the ellipse has dimensions $D_{25} \times d_{25}$ and is oriented along the optical major axis, $PA = 147^\circ$.

Fig. 13.— Position-velocity cuts obtained from the low resolution H I cube and the optical spectra for NGC 4772. Top panels: H I position-velocity diagrams along two slices through the low resolution data cube. The top right panel shows the slice along the optical major axis, $PA = 147^\circ$, along which the optical spectra were obtained. In addition to the H I contours, the velocities derived from the optical stellar absorption lines (open squares) and ionized gas emission lines (filled circles) are also shown. The top right panel shows the slice at $PA = 57^\circ$, along the minor axis. The lower panels show the optical rotation curves derived for NGC 4772 along the major axis ($PA = 147^\circ$) and along the minor axis ($PA = 57^\circ$). Separate symbols show the rotation curves derived from the stellar absorption and gas components. Error bars are shown.

Fig. 14.— Results for NGC 5854. Upper left panel: R band image. The ellipse has dimensions $D_{25} \times d_{25}$ and is oriented along the optical major axis, $PA = 55^\circ$. Upper right panel: Results of the two-dimensional bulge-disk decomposition for a $n = 4$ bulge. The solid line traces the light distribution while the disk and bulge are shown as long-dashed and short-dashed lines respectively. Center right panel: Central channels of the H I single dish spectrum obtained at Arecibo by Magri (1994). The lower panels show the optical rotation curves derived for NGC 5854 along the major axis ($PA = 55^\circ$) and along the minor axis ($PA = 145^\circ$). Separate symbols show the rotation curves derived from the stellar absorption and gas components. Error bars are shown.

Fig. 15.— New imaging results for NGC 4138. Top left panel: B band image scaled to illustrate the inner bulge, disk and dustlane. The ellipse has dimensions $D_{25} \times d_{25}$ and is oriented along the optical major axis, $PA = 150^\circ$. Top right panel: Results of the two-dimensional bulge-disk decomposition for a $n = 2$ bulge. The solid line traces the light distribution while the disk and bulge are shown as long-dashed and short-dashed lines respectively. Center left panel: The inner H I contours superposed on a zoomed B band image. The contours are at levels of 20 to 95×10^{19} in steps of 15×10^{19} atoms cm^{-2} . Center right panel: Similar display of the H I superposed on the H- α image. Bottom left panel: H I column density contours superposed on the V band image. The H I map shows contours at levels of 3.4, 6.9 and 13.8×10^{19} atoms cm^{-2} plus contours from 20 to 95×10^{19} atoms cm^{-2} in steps of 15 atoms cm^{-2} . For further details, see Jore *et al.* (1996). The ellipse is the same as in the upper left panel. Bottom right: Identical H I column density contours as the bottom left panel, here also shown as grayscale to emphasize the patchiness of the outer H I distribution.

This figure "Haynes.fig1.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0004340v1>

This figure "Haynes.fig2.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0004340v1>

This figure "Haynes.fig3.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0004340v1>

This figure "Haynes.fig4.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0004340v1>

This figure "Haynes.fig5.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0004340v1>

This figure "Haynes.fig6.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0004340v1>

This figure "Haynes.fig7.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0004340v1>

This figure "Haynes.fig8.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0004340v1>

This figure "Haynes.fig9.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0004340v1>

This figure "Haynes.fig10.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0004340v1>

This figure "Haynes.fig11.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0004340v1>

This figure "Haynes.fig12.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0004340v1>

This figure "Haynes.fig13.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0004340v1>

This figure "Haynes.fig14.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0004340v1>

This figure "Haynes.fig15.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0004340v1>